



Client: Rivn Research
Accession #: 2609010650
Search Code: Rivn2609010650
Received: 9/1/2026
Reported: 9/4/2026
Lot: KLOW-6844

Sample Summary

Product:	KLOW BLEND 80mg	Purity:	99.73%
Identity:	Confirmed	Net Content:	GHK-Cu – 57.92 mg KPV – 12.86 mg BPC-157 – 12.16 mg Thymosin Beta-4 – 10.17 mg
Appearance:	Blue Lyophilized Powder		
Endotoxin Threshold:	Pass		
Microbial Analysis (PCR):	Pass		
Elemental Impurities (ICP-MS): As Cd Pb Hg	Pass		

Analytical Results

Test	Result
Identity (LC-MS)	GHK-Cu/KPV/Thymosin Beta-4/BPC-157
Purity (HPLC-UV)	99.73%
Elemental Impurities (ICP-MS):	Pass — Arsenic, Cadmium, Lead, and Mercury below reporting/specification limits
Net Content	GHK-Cu – 57.92 mg KPV – 12.86 mg BPC-157 – 12.16 mg Thymosin Beta-4 – 10.17 mg

Method: Endotoxin testing performed using Limulus Amebocyte Lysate assay in accordance with USP <85> under validated laboratory conditions.


Endotoxin Replicate 1:	Pass	Assay Sensitivity: ≤0.05 EU/mL
Endotoxin Replicate 2:	Pass	Assay Sensitivity: ≤0.05 EU/mL

Method: Microbial detection performed using validated polymerase chain reaction (PCR)-based assay targeting common microbial contaminants.


Microbial Analysis (PCR)	No Detectable Microbial DNA
---------------------------------	-----------------------------

Method: HPLC with UV detection coupled with mass spectrometry (LC-MS).

Chromatogram



Mass Confirmation



Alex Johnson

Principal Chemist

FreedomDiagnosticsTesting.com

Admin@FreedomDiagnostics.net

Proudly Owned and Operated in the USA



The peptide purity analysis reported here was conducted using LCMS/MS under standard laboratory conditions. This analysis is intended for informational purposes only and is specific to the sample(s) provided. The peptides tested are intended for research use only and are not approved for human or veterinary use, diagnostic, therapeutic, or clinical applications. Results should be interpreted by qualified professionals within the scope of the intended research. The accuracy and reliability of the test may be influenced by sample integrity, handling, and other experimental variables.